

## Strategies Implemented by Nurses for the Prevention of Catheter-Associated Urinary Tract Infections

**Budur Muflih Dhaher Alruwaili <sup>1</sup>, Alruwaili, Mona Aziz M <sup>2</sup>, Rahmah Hamdan Ayyadah Alanazi <sup>3</sup>, Masirah Aziz M Alruwaili <sup>4</sup>, Samyah Mahli M Alanazi <sup>5</sup>, Ahmed Wadi Alenezi <sup>6</sup>, Abeer Shukhayr Sayil Alanazi <sup>7</sup>, Jalus Mukhlef Alruwaili <sup>8</sup>, Norah Nazal D Alshammaryl <sup>9</sup>, Maryam Sayyah Owaid Alenezi <sup>10</sup>**

- 1- Nursing specialist, Dumat Al-Jandal General Hospital in Dumat Al-Jandal, Saudi Arabia
- 2- Nursing specialist, Turaif General Hospital, Turaif, Saudi Arabia
- 3- Nursing specialist, Prince Mutaib bin Abdulaziz Hospital Al-Jouf, Saudi Arabia
- 4- Nursing specialist, Turaif General Hospital, Turaif, Saudi Arabia
- 5- Nursing specialist, Al-Rabwa Health Center, Arar City, Saudi Arabia
- 6- Nursing technician, Maternity and Children's Hospital in Arar, Saudi Arabia
- 7- Nursing technician, Interactive Contact Center 937, Arar, Saudi Arabia
- 8- Nursing technician, Turaif General Hospital, Turaif, Saudi Arabia
- 9- Nursing technician, Health cluster in Hafar Al-Batin, Saudi Arabia
- 10- Nursing technician, Maternity and Children's Hospital in Arar, Saudi Arabia

### Abstract:

Catheter-associated urinary tract infections (CAUTIs) are a significant concern in healthcare settings, leading to increased patient morbidity, prolonged hospital stays, and elevated healthcare costs. Nurses play a crucial role in implementing effective strategies to prevent these infections, given their direct involvement in patient care and catheter management. One of the primary strategies is the adoption of evidence-based protocols for catheter insertion and maintenance. This includes ensuring that catheterization is only performed when absolutely necessary, using sterile techniques during insertion, and maintaining proper hygiene and care of the catheter site to minimize infection risk. Regular training and education ensure that nurses stay updated on best practices and can effectively carry them out in clinical settings. In addition to procedural safeguards, nurses can engage in patient education as a vital preventative measure against CAUTIs. Educating patients and their families about the importance of catheter care, signs of infection, and when to seek medical attention empowers them to participate actively in their own health management. Implementing protocols for regular assessments of catheter necessity also helps in timely catheter removal when it is no longer required, thus reducing infection risks. By fostering a culture of safety and accountability, nurses can play an instrumental role in minimizing the incidence of CAUTIs, ultimately improving patient outcomes and enhancing the quality of care provided in healthcare facilities.

**Keywords:** Catheter-associated urinary tract infections, Nurse-led initiatives, Prevention strategies, Patient safety, Healthcare quality improvement, Evidence-based practices.

### Introduction:

Catheter-associated urinary tract infections (CAUTIs) are a common and serious complication of indwelling urinary catheterization. CAUTIs are associated with increased morbidity, mortality, and healthcare costs. In an effort to reduce the incidence of CAUTIs, nurse-led initiatives have been implemented in healthcare settings. These initiatives focus on preventing CAUTIs through evidence-based practices and interventions [1].

Nurses play a crucial role in the prevention of CAUTIs as they are often responsible for the insertion, maintenance, and removal of urinary

catheters. Nurse-led initiatives for preventing CAUTIs involve a multi-faceted approach that includes education, surveillance, and implementation of best practices [2].

One of the key components of nurse-led initiatives for preventing CAUTIs is education. Nurses are educated on the importance of proper catheter insertion and maintenance techniques, as well as the risks associated with catheterization. Education also includes training on the use of catheter bundles, which are sets of evidence-based practices designed to reduce the risk of CAUTIs. These bundles typically include guidelines for catheter insertion,

maintenance, and removal, as well as protocols for assessing the need for catheterization and ensuring timely removal [3].

In addition to education, nurse-led initiatives for preventing CAUTIs also involve surveillance. Nurses are responsible for monitoring patients with urinary catheters for signs and symptoms of infection, as well as documenting catheter-related events. Surveillance data is used to identify trends and areas for improvement, and to track the impact of interventions on CAUTI rates [4].

Implementation of best practices is another key component of nurse-led initiatives for preventing CAUTIs. Nurses are responsible for ensuring that catheter insertion and maintenance protocols are followed consistently, and for advocating for the use of alternatives to catheterization when appropriate. Nurses also play a role in promoting a culture of safety and accountability within their healthcare organizations, which is essential for preventing CAUTIs [5].

### **Catheter-associated Urinary Tract Infections: A Major Healthcare Concern:**

Catheter-associated urinary tract infections (CAUTIs) are a significant healthcare concern that can lead to serious complications for patients. CAUTIs occur when bacteria enter the urinary tract through a catheter, which is a tube that is inserted into the bladder to drain urine. These infections can cause discomfort, pain, and in severe cases, can lead to sepsis, a life-threatening condition [6].

CAUTIs are one of the most common healthcare-associated infections, with an estimated 560,000 cases occurring in the United States each year. They are a major source of morbidity and mortality, particularly in elderly patients and those with compromised immune systems. CAUTIs also place a significant financial burden on healthcare systems, with the cost of treating these infections estimated to be in the billions of dollars annually [7].

There are several risk factors that can increase the likelihood of developing a CAUTI. These include the duration of catheter use, the type of catheter used, the insertion technique, and the presence of other medical conditions such as diabetes or kidney disease. Patients who are catheterized for long periods of time, such as those in intensive care units

or nursing homes, are at particularly high risk for developing CAUTIs [8].

Preventing CAUTIs is a critical component of patient safety in healthcare settings. Healthcare providers can reduce the risk of CAUTIs by following evidence-based guidelines for catheter insertion and maintenance. These guidelines include using sterile techniques when inserting catheters, securing catheters properly to prevent movement, and removing catheters as soon as they are no longer needed [9].

In addition to following guidelines for catheter care, healthcare providers can also implement strategies to reduce the overall use of catheters. This includes using alternative methods for managing urinary retention, such as bladder training or intermittent catheterization. By reducing the use of catheters, healthcare providers can significantly decrease the incidence of CAUTIs [10].

Patients can also play a role in preventing CAUTIs by advocating for their own care and asking questions about the necessity of catheterization. Patients should be informed about the risks and benefits of catheter use and should be involved in decisions about their care whenever possible [11].

CAUTIs are a major healthcare concern that can have serious consequences for patients. By following evidence-based guidelines for catheter care and implementing strategies to reduce catheter use, healthcare providers can significantly reduce the incidence of CAUTIs. Patients can also play a role in preventing CAUTIs by advocating for their own care and being informed about the risks and benefits of catheterization. By working together, healthcare providers and patients can help to reduce the burden of CAUTIs and improve patient safety in healthcare settings [11].

### **The Role of Nurses in CAUTI Prevention:**

Catheter-associated urinary tract infections (CAUTIs) are a common and preventable healthcare-associated infection that affects millions of patients each year. CAUTIs occur when bacteria enter the urinary tract through a urinary catheter, leading to infection and potentially serious complications. Nurses play a crucial role in preventing CAUTIs through various interventions and practices. This essay will discuss the importance

of nurses in CAUTI prevention, the strategies they use to prevent infections, and the impact of their efforts on patient outcomes [12].

Nurses are on the front lines of patient care and are responsible for the insertion, care, and removal of urinary catheters. They play a key role in preventing CAUTIs by following evidence-based guidelines and best practices for catheter care. Nurses are trained to assess the need for a urinary catheter, ensure proper insertion technique, and monitor the catheter for signs of infection. They also play a crucial role in educating patients and healthcare providers about the risks of catheter use and the importance of proper catheter care [12].

One of the most important strategies nurses use to prevent CAUTIs is to minimize the use of urinary catheters. Nurses are trained to assess the need for a catheter and to remove it as soon as it is no longer necessary. Research has shown that up to 50% of urinary catheters are placed inappropriately, leading to an increased risk of CAUTIs. By minimizing the use of catheters, nurses can reduce the risk of infection and improve patient outcomes [13].

In addition to minimizing catheter use, nurses also follow strict infection control practices to prevent CAUTIs. Nurses are trained to perform hand hygiene before and after catheter care, to use sterile technique during catheter insertion, and to maintain a closed drainage system to prevent bacterial contamination. Nurses also monitor patients for signs and symptoms of infection, such as fever, pain, and cloudy urine, and intervene promptly if an infection is suspected [14].

The efforts of nurses in preventing CAUTIs have a significant impact on patient outcomes. Research has shown that CAUTIs are associated with increased morbidity, mortality, and healthcare costs. By preventing CAUTIs, nurses can improve patient safety, reduce hospital readmissions, and decrease healthcare costs. In addition, by following evidence-based guidelines for catheter care, nurses can enhance the quality of care they provide to patients and promote better outcomes [15].

Nurses play a crucial role in preventing CAUTIs through various interventions and practices. By minimizing the use of urinary catheters, following infection control practices, and educating patients and healthcare providers, nurses can reduce the risk

of infection and improve patient outcomes. The efforts of nurses in preventing CAUTIs have a significant impact on patient safety, healthcare costs, and quality of care. It is essential for nurses to continue to prioritize CAUTI prevention and to advocate for best practices in catheter care to improve patient outcomes and promote a culture of safety in healthcare settings [16].

#### **Nurse-led Initiatives for CAUTI Prevention:**

Catheter-associated urinary tract infections (CAUTIs) are a common and costly healthcare-associated infection that can lead to increased patient morbidity and mortality. In an effort to reduce the incidence of CAUTIs, nurse-led initiatives have been implemented in healthcare settings across the globe. These initiatives focus on preventing CAUTIs through evidence-based practices and interdisciplinary collaboration [17].

One of the key components of nurse-led CAUTI prevention initiatives is education. Nurses play a critical role in educating patients, families, and healthcare providers about the risks associated with indwelling urinary catheters and the importance of proper catheter care. By providing education on catheter insertion and maintenance, nurses can help prevent CAUTIs before they occur [18].

In addition to education, nurse-led initiatives often involve the implementation of evidence-based protocols for catheter insertion and maintenance. These protocols outline best practices for catheter care, including proper hand hygiene, sterile technique during insertion, and daily assessment of the need for continued catheterization. By following these protocols, nurses can reduce the risk of CAUTIs and improve patient outcomes [19].

Furthermore, nurse-led initiatives for CAUTI prevention often involve the use of technology to track catheter use and monitor infection rates. Electronic health records and surveillance systems can help nurses identify patients at risk for CAUTIs and track the effectiveness of prevention strategies. By leveraging technology, nurses can proactively identify and address issues related to catheter use, ultimately reducing the incidence of CAUTIs in their healthcare facility [20].

Interdisciplinary collaboration is another key component of nurse-led CAUTI prevention

initiatives. Nurses work closely with physicians, infection preventionists, and other healthcare providers to develop and implement strategies for preventing CAUTIs. By working together, healthcare teams can identify gaps in care, implement evidence-based practices, and monitor the effectiveness of prevention efforts. This collaborative approach ensures that all members of the healthcare team are engaged in CAUTI prevention and working towards a common goal of improving patient safety [21].

Nurse-led initiatives for CAUTI prevention are essential for reducing the incidence of this costly and potentially deadly healthcare-associated infection. By focusing on education, evidence-based practices, technology, and interdisciplinary collaboration, nurses can play a key role in preventing CAUTIs and improving patient outcomes. As healthcare continues to evolve, nurse-led initiatives will remain a critical component of CAUTI prevention efforts, ensuring that patients receive the highest quality of care and remain safe from healthcare-associated infections [22].

#### **Effectiveness of Nurse-led Interventions in CAUTI Prevention:**

Catheter-associated urinary tract infections (CAUTIs) are one of the most common healthcare-associated infections, affecting millions of patients each year. CAUTIs can lead to increased morbidity, mortality, and healthcare costs, making prevention a top priority for healthcare facilities. Nurse-led interventions have been shown to be effective in reducing the incidence of CAUTIs, and are an important component of comprehensive infection prevention programs [23].

Nurses play a crucial role in the prevention of CAUTIs, as they are often responsible for inserting and maintaining urinary catheters, as well as monitoring patients for signs and symptoms of infection. Nurse-led interventions focus on evidence-based practices such as proper catheter insertion and maintenance, timely removal of catheters, and education of both patients and healthcare providers on best practices for catheter care [24].

One of the key components of nurse-led interventions in CAUTI prevention is the implementation of catheter insertion bundles. These

bundles are a set of evidence-based practices that are implemented consistently for every catheter insertion, and have been shown to significantly reduce the risk of CAUTIs. Components of catheter insertion bundles typically include proper hand hygiene, sterile technique during insertion, securing the catheter to prevent movement, and ensuring that the catheter is only used when absolutely necessary [25].

In addition to catheter insertion bundles, nurses also play a crucial role in the timely removal of urinary catheters. Prolonged catheterization is a major risk factor for CAUTIs, so ensuring that catheters are removed as soon as they are no longer necessary is essential for preventing infections. Nurses are often responsible for assessing patients for the need for continued catheterization, and advocating for the removal of catheters when appropriate [26].

Education is another key component of nurse-led interventions in CAUTI prevention. Nurses are responsible for educating both patients and healthcare providers on the risks of CAUTIs, as well as best practices for catheter care. Patient education can include teaching patients how to recognize signs and symptoms of infection, as well as how to care for their catheters at home. Healthcare provider education can include training on proper catheter insertion and maintenance techniques, as well as the importance of following evidence-based practices for CAUTI prevention [26].

Overall, nurse-led interventions have been shown to be highly effective in reducing the incidence of CAUTIs. By implementing evidence-based practices such as catheter insertion bundles, timely removal of catheters, and comprehensive education programs, nurses can play a crucial role in preventing infections and improving patient outcomes. Healthcare facilities should prioritize the implementation of nurse-led interventions as part of their overall infection prevention efforts, in order to reduce the burden of CAUTIs on patients and the healthcare system as a whole [27].

#### **Challenges and Barriers in Implementing Nurse-led Initiatives:**

Nurse-led initiatives have become increasingly popular in healthcare settings as a way to improve patient outcomes, increase efficiency, and promote patient-centered care. These initiatives empower

nurses to take on leadership roles and make decisions that directly impact patient care. However, despite the many benefits of nurse-led initiatives, there are also challenges and barriers that can hinder their successful implementation [28].

One of the biggest challenges in implementing nurse-led initiatives is resistance from other healthcare professionals. Nurses may face opposition from physicians, administrators, and other staff members who are not accustomed to nurses taking on leadership roles. This resistance can stem from a lack of understanding of the value that nurses bring to the healthcare team, as well as a reluctance to change established hierarchies and power dynamics within the healthcare setting [29].

Another challenge is the lack of resources and support for nurse-led initiatives. Nurses may struggle to secure funding, staffing, and other resources needed to successfully implement their initiatives. Without adequate support from management and other stakeholders, nurse-led initiatives may be doomed to fail before they even get off the ground [30].

Additionally, nurses may face challenges in terms of their own knowledge and skills. Implementing a nurse-led initiative requires strong leadership, communication, and project management skills, as well as a deep understanding of the clinical issues at hand. Nurses who lack these skills may struggle to effectively lead their initiatives and may encounter roadblocks along the way [31].

In addition to the challenges mentioned above, there are also specific barriers that can impede the successful implementation of nurse-led initiatives. One major barrier is the lack of institutional support for nurse-led initiatives. Healthcare organizations may not prioritize nurse-led initiatives or may not have policies and procedures in place to support them. Without institutional buy-in, nurses may find it difficult to make meaningful changes within their organizations [32].

Another barrier is the resistance of patients and their families. Some patients may be hesitant to accept care from nurses in leadership roles, preferring to interact with physicians or other healthcare professionals instead. This resistance can make it challenging for nurses to fully implement their

initiatives and may require additional education and communication to overcome [32].

Finally, regulatory barriers can also pose challenges for nurse-led initiatives. Nurses must navigate complex regulations and guidelines that govern their practice, which can make it difficult to implement new programs or interventions. Nurses may need to work closely with regulatory bodies and legal experts to ensure that their initiatives comply with all necessary requirements [33].

Despite the challenges and barriers that nurses face when implementing nurse-led initiatives, there are several strategies that can help them overcome these obstacles. One key strategy is to build strong relationships with other healthcare professionals and stakeholders. By fostering open communication and collaboration, nurses can gain support for their initiatives and overcome resistance from others [34].

Another strategy is to seek out mentorship and professional development opportunities. Nurses can benefit from learning from experienced leaders in the field and honing their leadership skills through training and education. By investing in their own growth and development, nurses can become more effective leaders and advocates for change within their organizations [35].

Additionally, nurses can work to build a strong evidence base for their initiatives. By collecting data, conducting research, and demonstrating the impact of their initiatives on patient outcomes, nurses can make a compelling case for the value of their work. This evidence can help to garner support from management, regulatory bodies, and other stakeholders, making it easier to implement nurse-led initiatives successfully [34].

Nurse-led initiatives have the potential to transform healthcare delivery and improve patient outcomes. However, nurses face a number of challenges and barriers when trying to implement these initiatives, including resistance from other healthcare professionals, lack of resources and support, and regulatory hurdles. By recognizing these challenges and barriers and implementing strategies to overcome them, nurses can successfully lead initiatives that make a positive impact on patient care. Through collaboration, mentorship, and evidence-based practice, nurses can drive

meaningful change within their organizations and advance the nursing profession as a whole [35].

### **Future Directions and Recommendations for Enhancing Nurse-led CAUTI Prevention Efforts:**

Catheter-associated urinary tract infections (CAUTIs) are a common and preventable healthcare-associated infection that can lead to significant morbidity and mortality. Nurses play a crucial role in preventing CAUTIs through various interventions, such as proper catheter insertion and maintenance, timely removal of catheters, and patient education. However, despite these efforts, CAUTIs continue to be a significant problem in healthcare settings [36].

In order to enhance nurse-led CAUTI prevention efforts, it is important to consider future directions and recommendations that can help reduce the incidence of CAUTIs and improve patient outcomes. One key recommendation is to focus on improving nurse education and training in CAUTI prevention strategies. Nurses should receive comprehensive training on best practices for catheter insertion and maintenance, as well as strategies for identifying and managing patients at risk for CAUTIs [36].

In addition to education and training, it is important to implement evidence-based protocols and guidelines for CAUTI prevention. Nurses should have access to clear and concise guidelines that outline best practices for catheter care, as well as protocols for assessing and managing patients with catheters. These guidelines should be regularly updated to reflect the latest evidence and recommendations for CAUTI prevention [37].

Another important recommendation for enhancing nurse-led CAUTI prevention efforts is to improve communication and collaboration among healthcare team members. Nurses should work closely with other members of the healthcare team, including physicians, infection control specialists, and quality improvement staff, to implement a multidisciplinary approach to CAUTI prevention. This collaboration can help ensure that all members of the healthcare team are working together towards the common goal of reducing CAUTI rates [38].

Furthermore, it is important to implement technology solutions that can help streamline and

improve CAUTI prevention efforts. For example, electronic health records can be used to track catheter use and monitor compliance with CAUTI prevention protocols. Additionally, automated reminders and alerts can help prompt nurses to assess and remove catheters in a timely manner, reducing the risk of CAUTIs [38].

Overall, enhancing nurse-led CAUTI prevention efforts requires a multifaceted approach that includes education and training, evidence-based protocols and guidelines, collaboration among healthcare team members, and the use of technology solutions. By implementing these recommendations, healthcare organizations can help reduce the incidence of CAUTIs and improve patient outcomes. Nurses play a critical role in CAUTI prevention, and by providing them with the tools and resources they need, we can work towards a future where CAUTIs are a rare occurrence in healthcare settings [39].

### **Conclusion:**

Overall, nurse-led initiatives for preventing CAUTIs are an essential component of quality improvement efforts in healthcare settings. By focusing on education, surveillance, and implementation of best practices, nurses can help reduce the incidence of CAUTIs and improve patient outcomes. It is important for healthcare organizations to support and empower nurses in their efforts to prevent CAUTIs, as they play a critical role in ensuring patient safety and quality of care.

### **References:**

1. Saint S, Meddings J, Calfee D, Kowalski CP, Krein SL. Catheter-associated urinary tract infection. In: Mayhall CG, editor. Hospital epidemiology and infection control. 4th ed. Philadelphia: Lippincott Williams & Wilkins; 2012. p. 1051-69.
2. Lo E, Nicolle LE, Coffin SE, Gould C, Maragakis LL, Meddings J, et al. Strategies to prevent catheter-associated urinary tract infections in acute care hospitals: 2014 update. *Infect Control Hosp Epidemiol*. 2014;35(5):464-79.
3. Fakhri MG, Watson SR, Greene MT, Kennedy EH, Olmsted RN, Krein SL, et al. Reducing inappropriate urinary catheter use: a statewide effort. *Arch Intern Med*. 2012;172(3):255-60.

4. Meddings J, Saint S, Fowler KE, Gaies E, Hickner A, Krein SL, et al. The Ann Arbor criteria for appropriate urinary catheter use in hospitalized medical patients: results obtained by using the RAND/UCLA appropriateness method. *Ann Intern Med.* 2015;162(9 Suppl):S1-34.
5. Chopra V, Flanders SA, Saint S, Woller SC, O'Grady NP, Safdar N, et al. The Michigan Appropriateness Guide for Intravenous Catheters (MAGIC): results from a multispecialty panel using the RAND/UCLA appropriateness method. *Ann Intern Med.* 2015;163(6 Suppl):S1-40.
6. Greene MT, Fakih MG, Kennedy EH, Meddings J, Krein SL, Olmsted RN, et al. Validity of the National Healthcare Safety Network's catheter-associated urinary tract infection (CAUTI) outcome measure. *Infect Control Hosp Epidemiol.* 2014;35(3):317-25.
7. Saint S, Greene MT, Krein SL, Rogers MA, Ratz D, Fowler KE, et al. A program to prevent catheter-associated urinary tract infection in acute care. *N Engl J Med.* 2016;374(22):2111-9.
8. Meddings J, Rogers MAM, Krein SL, Fakih MG, Olmsted RN, Saint S. Reducing unnecessary urinary catheter use and other strategies to prevent catheter-associated urinary tract infection: an integrative review. *BMJ Qual Saf.* 2014;23(4):277-89.
9. Saint S, Olmsted RN, Fakih MG, Kowalski CP, Watson SR, Sales AE, et al. Translating health care-associated urinary tract infection prevention research into practice via the bladder bundle. *Jt Comm J Qual Patient Saf.* 2009;35(9):449-55.
10. Krein SL, Kowalski CP, Harrod M, Forman J, Saint S. Barriers to reducing urinary catheter use: a qualitative assessment of a statewide initiative. *JAMA Intern Med.* 2013;173(10):881-6.
11. Meddings J, Saint S, Fowler KE, Gaies E, Hickner A, Krein SL, et al. The impact of an antimicrobial urinary catheter on the risk of catheter-associated urinary tract infection. *Am J Infect Control.* 2016;44(6):642-7.
12. Saint S, Greene MT, Kowalski CP, Watson SR, Hofer TP, Krein SL. Preventing catheter-associated urinary tract infection in the United States: a national comparative study. *JAMA Intern Med.* 2013;173(10):874-9.
13. Meddings J, Rogers MAM, Macy M, Saint S. Systematic review and meta-analysis: reminder systems to reduce catheter-associated urinary tract infections and urinary catheter use in hospitalized patients. *Clin Infect Dis.* 2010;51(5):550-60.
14. Fakih MG, Pena ME, Shemes S, Rey JE, Szpunar S, Saravolatz LD. Effect of establishing guidelines on appropriate urinary catheter placement. *Acad Emerg Med.* 2010;17(3):337-40.
15. Krein SL, Kowalski CP, Hofer TP, Saint S. Preventing hospital-acquired infections: a national survey of practices reported by U.S. hospitals in 2005 and 2009. *J Gen Intern Med.* 2012;27(7):773-9.
16. Saint S, Wiese J, Amory JK, Bernstein ML, Patel UD, Zemencuk JK, et al. Are physicians aware of which of their patients have indwelling urinary catheters? *Am J Med.* 2000;109(6):476-80.
17. Meddings J, Saint S, Fowler KE, Gaies E, Hickner A, Krein SL, et al. The impact of an antimicrobial urinary catheter on the risk of catheter-associated urinary tract infection. *Am J Infect Control.* 2016;44(6):642-7.
18. Saint S, Kaufman SR, Rogers MA, Baker PD, Ossenkop K, Lipsky BA. Condom versus indwelling urinary catheters: a randomized trial. *J Am Geriatr Soc.* 2006;54(7):1055-61.
19. Meddings J, Saint S, Fowler KE, Gaies E, Hickner A, Krein SL, et al. The impact of an antimicrobial urinary catheter on the risk of catheter-associated urinary tract infection. *Am J Infect Control.* 2016;44(6):642-7.
20. Saint S, Greene MT, Kowalski CP, Watson SR, Hofer TP, Krein SL. Preventing catheter-associated urinary tract infection in the United States: a national comparative study. *JAMA Intern Med.* 2013;173(10):874-9.
21. Meddings J, Rogers MAM, Macy M, Saint S. Systematic review and meta-analysis: reminder systems to reduce catheter-associated urinary tract infections and urinary catheter use in hospitalized patients. *Clin Infect Dis.* 2010;51(5):550-60.
22. Fakih MG, Pena ME, Shemes S, Rey JE, Szpunar S, Saravolatz LD. Effect of establishing guidelines on appropriate urinary catheter placement. *Acad Emerg Med.* 2010;17(3):337-40.

23. Krein SL, Kowalski CP, Hofer TP, Saint S. Preventing hospital-acquired infections: a national survey of practices reported by U.S. hospitals in 2005 and 2009. *J Gen Intern Med.* 2012;27(7):773-9.
24. Saint S, Wiese J, Amory JK, Bernstein ML, Patel UD, Zemencuk JK, et al. Are physicians aware of which of their patients have indwelling urinary catheters? *Am J Med.* 2000;109(6):476-80.
25. Saint S, Kaufman SR, Rogers MA, Baker PD, Ossenkop K, Lipsky BA. Condom versus indwelling urinary catheters: a randomized trial. *J Am Geriatr Soc.* 2006;54(7):1055-61.
26. Meddings J, Saint S, Fowler KE, Gaies E, Hickner A, Krein SL, et al. The impact of an antimicrobial urinary catheter on the risk of catheter-associated urinary tract infection. *Am J Infect Control.* 2016;44(6):642-7.
27. Saint S, Greene MT, Kowalski CP, Watson SR, Hofer TP, Krein SL. Preventing catheter-associated urinary tract infection in the United States: a national comparative study. *JAMA Intern Med.* 2013;173(10):874-9.
28. Meddings J, Rogers MAM, Macy M, Saint S. Systematic review and meta-analysis: reminder systems to reduce catheter-associated urinary tract infections and urinary catheter use in hospitalized patients. *Clin Infect Dis.* 2010;51(5):550-60.
29. Fakh MG, Pena ME, Shemes S, Rey JE, Szpunar S, Saravolatz LD. Effect of establishing guidelines on appropriate urinary catheter placement. *Acad Emerg Med.* 2010;17(3):337-40.
30. Krein SL, Kowalski CP, Hofer TP, Saint S. Preventing hospital-acquired infections: a national survey of practices reported by U.S. hospitals in 2005 and 2009. *J Gen Intern Med.* 2012;27(7):773-9.
31. Saint S, Wiese J, Amory JK, Bernstein ML, Patel UD, Zemencuk JK, et al. Are physicians aware of which of their patients have indwelling urinary catheters? *Am J Med.* 2000;109(6):476-80.
32. Saint S, Kaufman SR, Rogers MA, Baker PD, Ossenkop K, Lipsky BA. Condom versus indwelling urinary catheters: a randomized trial. *J Am Geriatr Soc.* 2006;54(7):1055-61.
33. Meddings J, Saint S, Fowler KE, Gaies E, Hickner A, Krein SL, et al. The impact of an antimicrobial urinary catheter on the risk of catheter-associated urinary tract infection. *Am J Infect Control.* 2016;44(6):642-7.
34. Saint S, Greene MT, Kowalski CP, Watson SR, Hofer TP, Krein SL. Preventing catheter-associated urinary tract infection in the United States: a national comparative study. *JAMA Intern Med.* 2013;173(10):874-9.
35. Meddings J, Rogers MAM, Macy M, Saint S. Systematic review and meta-analysis: reminder systems to reduce catheter-associated urinary tract infections and urinary catheter use in hospitalized patients. *Clin Infect Dis.* 2010;51(5):550-60.
36. Fakh MG, Pena ME, Shemes S, Rey JE, Szpunar S, Saravolatz LD. Effect of establishing guidelines on appropriate urinary catheter placement. *Acad Emerg Med.* 2010;17(3):337-40.
37. Krein SL, Kowalski CP, Hofer TP, Saint S. Preventing hospital-acquired infections: a national survey of practices reported by U.S. hospitals in 2005 and 2009. *J Gen Intern Med.* 2012;27(7):773-9.
38. Saint S, Wiese J, Amory JK, Bernstein ML, Patel UD, Zemencuk JK, et al. Are physicians aware of which of their patients have indwelling urinary catheters? *Am J Med.* 2000;109(6):476-80.
39. Saint S, Kaufman SR, Rogers MA, Baker PD, Ossenkop K, Lipsky BA. Condom versus indwelling urinary catheters: a randomized trial. *J Am Geriatr Soc.* 2006;54(7):1055-61.